


This article was downloaded by: [Gérvas, Juan]

On: 16 December 2009

Access details: Access Details: [subscription number 917455327]

Publisher Informa Healthcare

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



The European Journal of General Practice

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t749989318>

Clinical prevention: patients' fear and the doctor's guilt

Juan Gérvas ^a; Iona Heath ^a; Antonio Durán ^a; Joan Gené The Members of the Seminar of Primary Health Innovation 2008 (Seminario de Innovación en atención primaria 2008) ^a

^a Equipo CESCA, Travesía de la Playa, 3, 28730 Buitrago del Lozoya, Madrid, Spain

Online publication date: 05 December 2009

To cite this Article Gérvas, Juan, Heath, Iona, Durán, Antonio and Gené The Members of the Seminar of Primary Health Innovation 2008 (Seminario de Innovación en atención primaria 2008), Joan(2009) 'Clinical prevention: patients' fear and the doctor's guilt', The European Journal of General Practice, 15: 3, 122 – 124

To link to this Article: DOI: 10.3109/13814780903242481

URL: <http://dx.doi.org/10.3109/13814780903242481>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

SEMINAR REPORT

Clinical prevention: patients' fear and the doctor's guilt

JUAN GÉRVAS, IONA HEATH, ANTONIO DURÁN, JOAN GENÉ, THE MEMBERS OF THE SEMINAR OF PRIMARY HEALTH INNOVATION 2008 (SEMINARIO DE INNOVACIÓN EN ATENCIÓN PRIMARIA 2008)

Equipo CESCA, Travesía de la Playa, 3, 28730 Buitrago del Lozoya, Madrid, Spain

Introduction

Contrary to its apparently straightforward concept, the word “prevention” has become a sort of “mantra” whose meaning is difficult to establish, at least in relation to health services (1). In its application in the wide field of health, prevention has a different significance depending on whether it is related to populations or to patients (2,3).

Clinical prevention is the activity which anticipates and tries to avoid and reduce harm, generally offered to patients at the doctor's office by health professionals (4). With the introduction of the risk factor concept, the clinical prevention field has substantially broadened, practically without limits, as is clearly demonstrated, for example, by the numerous cardiovascular risk factors currently described (5). A risk factor era has emerged as the result of the growing predominance of clinical activity and public health programmes intended to minimize the risks for certain diseases, especially cardiovascular disease and cancer (6). Furthermore, the appearance of the concept of “pre-illness” increases the scope of prevention, as the goal becomes to anticipate and treat situations which could result in established disease. A good example of this is pre-diabetes (7).

Prevention has some unquestionable successes. However, the extension of its influence has brought about proposals for activities which are not very efficient in clinical prevention. It is not clear why doctors and patients have accepted such clinical prevention, considering the huge impact it has on their jobs and daily life. There are two main issues involved: feelings of fear (among patients) and feelings of guilt (among doctors).

Patient fear

Patients go to the doctor with a number of expectations and fears. For example, a patient may be worried about a surprising number of differential diagnoses when he or she goes to the doctor for a “simple” acute cough (8). Patients fear for their lives, they worry about suffering from diseases and/or becoming seriously disabled (9).

Fear makes patients feel vulnerable, as they put themselves in the hands of doctors, expecting a solution to everything. It also gives rise to unfounded expectations of avoiding disease and death due to *zero risk*, and systematic control of all risk factors (10).

Excessively high expectations among patients bring with them the desire for perfect health, which paradoxically creates the feeling of having poor health (11). Furthermore, patients consent to medical interventions when the benefit is doubtful or marginal. With regards to screening, the inclusion of routine breast ultrasound may be used as an example. This test decreases the positive predictive value of the recommendation of breast surgery from 23% to 11% (12). In the desire to be in perfect health, in order to meet all expectations and eliminate all fears, prevention could do more harm than good (13).

In order to avoid the harm of excessive prevention activity, the doctor should consider the fears and expectations of the patient, within the consultation, and provide appropriate answers.

Doctor's guilt

Disease was once related to sin, whereas nowadays it tends to be associated with a “possible error in clinical

prevention". Because of this, there still exists some feeling of blame, currently transformed into the doctor's feelings of guilt regarding the patient, for example, with a myocardial infarction. The doctor immediately goes through the patient's clinical record and detects the *missed opportunities* for prevention.

Doctors feel guilty in the face of disease and suffering. It is not sufficient to be a healer; it seems that modern times demand that you should also be a scientist and apply the guidelines and norms which ensure good results. The guidelines, norms, and protocols stop being an aid to decision making and become instructions which must be followed.

Clinical practice becomes prescriptive and ignores the way patients think about and experience their own lives and illnesses. It is said that "there are no illnesses, just ill people", but it is difficult to resist the pressures of normative "standards of care", even knowing that the sick person himself must remain in charge of his own personal evolution.

We cannot ignore the obvious—the indisputable law of epidemiology—that "all that lives, dies" (14). We should be honest with patients and let them know that there are cases of clinical prevention which simply change the cause of death, and do not even delay the process or allow one to die with dignity. For example, in the case of cardiovascular prevention in the elderly, life is not extended but the cause of death may be different (dementia and cancer) (15).

Doctors should act responsibly but should not be made to feel guilty for arbitrary and contingent events. This increases unhappiness in the doctor as a professional, but it is impossible to respond to the excessively high expectations of prevention without limits (10,16). Currently, medical training is still based on the illness and its diagnosis, which contributes to the frustration caused by the contrast between training and daily clinical practice (16,17).

Conclusion

Clinical prevention should not be allowed to evolve into a categorical imperative in which the end justifies the means, which fuels patients' fears and doctor's guilt, and which endorses clinical intervention with a certain sense of righteousness in the search for good (18). Clinical prevention is an important activity which nearly always has a place in the doctor-patient consultation (19). However, clinical prevention has its limits (20). Above all, the activities of clinical prevention should not be introduced/maintained as a result of feelings of fear (in the patient) or guilt (in the doctor).

Funding body

The Seminars on Innovation in Primary Care (Innovación en Atención Primaria) are organized by the Foundation of Health Sciences (Fundación Ciencias de la Salud) and the Foundation for Training of the Spanish Medical College Organization (Fundación para la Formación, Organización Médica Colegial) with the cooperation of the Institute for Research in Primary Care Jordi Gol (IDIAP Jordi Gol) and the sponsorship of the Minister of Health and GSK. We thank IDIAP Jordi Gol for financial help in preparing the English version of the manuscript.

Competing interests

The authors have stated that there are none.

Acknowledgements

This paper is a report on the 1st Seminar on Innovation in Primary Care 2008 (Madrid, 7th March 2008) held under the auspices of the Fundación Ciencias de la Salud and the Fundación para la Formación de la Organización Médica Colegial, and sponsored by Glaxo-Smith-Kline.

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

References

1. Starfield B, Hyde J, Gervas J, Heath I. The concept of prevention: a good idea gone astray? *J Epidemiol Community Health* 2008;62:580–3.
2. Rose G. Sick individuals and sick populations. *Int J Epidemiol* 1985;14:32–8.
3. Charlton BG. A critique of Geoffrey Rose's 'population strategy' for preventive medicine. *J R Soc Med* 1995;88:607–10.
4. Gray JA. New concepts in screening. *Br J Gen Pract* 2004;54:292–8.
5. Brotman DJ, Walker E, Lauer MS, O'Brien RG. In search of fewer independent risk factors. *Arch Intern Med* 2005;165:138–45.
6. Stoltenberg C. Merging genetics and epidemiology: what is in it for public health? *Scand J Public Health* 2005;33:1–3.
7. Nissen SE, Wolski K. Effect of rosiglitazone on the risk of myocardial infarction and death from cardiovascular causes. *N Engl J Med* 2007;356:2457–71.
8. Bergh KD. The patient's differential diagnosis. Unpredictable concerns in visits for acute cough. *J Fam Pract* 1998;46:153–8.
9. WONCA International Classification Committee. International classification of primary care, Oxford: Oxford University Press; 1998.
10. Fugelli P. The zero-vision: potential side effects of communicating health perfection and zero risk. *Patient Educ Couns* 2006;60:267–71.

11. Sen A. Health: perception versus observation. *BMJ* 2002;324:860–1.
12. Berg WA, Blume JD, Cormack JB, Mendelson EB, Lehrer D, Bohm-Velez M, et al. Combined screening with ultrasound and mammography vs mammography alone in women at elevated risk of breast cancer. *JAMA* 2008;299:2151–63.
13. Illich I. L'obsession de la santé parfaite. *Le Monde Diplomatique*. Available from: URL:www.monde-diplomatique.fr/1999/03/ILLICH/11802
14. Ebrahim S. Chronic diseases and calls to action. *Int J Epidemiol* 2008;37:225–30.
15. Mangin D, Sweeney K, Heath I. Preventive health care in elderly people needs rethinking. *BMJ* 2007;335:285–7.
16. Smith R. Why are doctors so unhappy? There are probably many causes, some of them deep. *BMJ* 2001;322:1073–4.
17. Peile E, Carter Y. Selecting and supporting contented doctors. *BMJ* 2005;330:269–70.
18. Smith GD. Reflections on the limitations to epidemiology. *J Clin Epidemiol* 2001;54:325–31.
19. Stott NC, Davis RH. The exceptional potential in each primary care consultation. *J R Coll Gen Pract* 1979;29:201–5.
20. Gervas J, Starfield B, Heath I. Caution in clinical prevention. *Lancet* 2008;372:1997–9.